

# CERTIFICATE OF ANALYSIS

# U-[13C34]-Fumonisin B2 in Acetonitrile/Water (50/50) LCMS grade

This document is designed, and the certified values and uncertainty are determined in accordance with ISO Guide 31, ISO Guide 34, ISO Guide 35, AOAC, and Eurachem/ CITAC Guides.

### Description of the Reference Material (RM)



#### Intended use of the RM:

For laboratory use for R&D purposes only. The main purpose of this material is for analytical instrument calibration (e. g. external calibration, standard addition). Not for drug, household or other uses.

#### Instruction for the correct use of the RM:

The vial should be stored in a dark place at 2.8°C. Before usage of the RM, allow the vial to warm to room temperature. The expiry date of this RM is based on the current knowledge and holds only for proper storage conditions in the originally closed vials? packages. Solutions prepared for calibration purpose should be protected from exposure to light. Discard solutions after us in accordance with appropriate saffer progulations for chemical substances.

### Hazardous situation:

H225 : Flammable liquid - Category 2 - Highly flammable liquid and vapour

H302 : Acute toxicity - Oral - Category 4 - Harmful if swallowed
H319 : Eye irritation - Category 2 - Causes serious eye irritation

In case of accident or if you feel unwell, seek medical advice immediately (show the label where possible). Avoid exposure. Wear suitable protective clothing

## Safety measures:

Special care must be taken when manipulating this standard. Avoid contact with eyes, skin and clothing. Avoid prolonged or repeated exposure. Use only in a chemical fune hood. Safety shower and eye bath must be near. In case of spills, cover and absorb with an inert dry material such as dry-time, sand or soda ash and place in an appropriate waste disosocial container.

Keep container tightly closed. Do not store in direct sunlight. Keep away from heat, sparks, flame and incompatible material. Storage area should be cool, dry and away from incompatible materials.

## Further information:

Further information is available in the MSDS provided along with this certificate. Final users should make their own investigations to determine the suitability of the information for their particular research purposes. In no event the supplier of this RM shall be held liable for any damage resulting from handling or from contact with the product.

### Traceability

The certified values are based on the results of analytical techniques previously used for purity assessment of solid mycotoxins. High purity material represents a practical realization of concentration units, through conversion of mass to molar quantity.



## Calculation of certified values and associated uncertainties

This calibrant is certified on solution preparation. Toxin is pipetted and diluted in Acetonitrile/Water (50/50) LCMS grade. Mass concentration calculation is based on certified concentration, purity and dilution step.

The pipet was calibrated with traceability to national and international standards (Dakks & ilac-MRA). All weights used for metrological control are connected to national and international standards. The weights are calibrated by an accredited laboratory.

$$C (\mu g/mL) = \frac{m \times P}{\nu}$$

Toxin	Source				Standard uncertainty
U-[13C34]-Fumonisin B2	Purity		100,00	%	0,00
	Liquid solution	concentration	3000,00	μg/mL	0,92
	Volumetry procedure	volume	0,17	mL	0,00
	Dilution	volume	50,00	mL	0,06
$Combined_a = \sqrt{\left(\frac{n_b}{p}\right)^2 + \left(\frac{n_{con}}{V_{con}}\right)^2 + \left(\frac{n_{rp}}{V_{co}}\right)^2 + \left(\frac{n_{rp}}{V_{c}}\right)^2 + \left(\frac{n_{rp}}{V_{c}}\right)^2}$					0,01
Concentration <sub>Parkin</sub> = Concentration nother V <sub>E1</sub> µg/mL					10,20
Total expanded uncertainty (using a coverage factor k=2)					0,13

Notes:

The purity of the mycotoxin used for this RM was determined by liquid chromatography.


Following the Guide to the Expression of Uncertainty in measurement (GUM) the expanded uncertainty of toxin level is obtained by multiplication with a coverage factor K for which 2 is usually chosen to obtain a confidence level of 95 %.

#### Carbon 13 calculation

Isotopic incorporation				
Compound	Isotopic distribution			
<sup>13</sup> C <sub>31</sub> Fumonisin B2	3,6%			
<sup>13</sup> C <sub>∞</sub> Fumonisin B2	10,2%			
<sup>13</sup> C <sub>33</sub> Fumonisin B2	28,9%			
<sup>13</sup> C <sub>34</sub> Fumonisin B2	57,3%			
Calculated isotopic incorporation ( 13 C/molecule)	98,4%			

The calculation are based on LC-MS/MS data

## Quality control





## References:

a-ISO GUIDE 31:2015, Reference Materials - Contents of certificates, labels and accompanying documentation.

b-ISO GUIDE 34:2009, General requirements for the competence of reference material producers

c-ISO GUIDE 35:2006, Reference materials - General and Statistical Principles.

d-ISO/IEC Guide 98-3:2008 Uncertainty of measurement-Part 3 : Guide to the expression of uncertainty in measurement (GUM:1995)

e-Eurachem/CITAC guide (2019), Traceability in Chemical Measurement. f-Eurachem/CITAC guide (2012), Quantifying Uncertainty in Analytical Measurement.

g-AOAC Official Method 970.44-1971 - Preparation of Standards for Mycotoxins.

# This document was computer generated and is valid without signature

Prepared by: CLERMONT Alexandre Quality Control

Date: 01-Jul-24